


IN THE CLAIMS

Please amend the claims as follows:

Claim 1 (Currently Amended): A production method of a pyridine compound
~~derivative~~ having a substituent at the 2-position of an aromatic heterocyclic structure, which
 is represented by the formula (III')




wherein

R^2 , R^3 , R^4 and R^5

are each a hydrogen atom, a halogen atom, an alkyl group
 optionally having substituent(s), an aryl group
 optionally having substituent(s), an alkoxy group
 optionally having substituent(s), an aryloxy group
 optionally having substituent(s), an acyloxy group
 optionally having substituent(s), an alkylthio group
 optionally having substituent(s), an arylthio group
 optionally having substituent(s), an arylthio group
 optionally having substituent(s), a protected amino
 group optionally having substituent(s), a nitro group, a
 cyano group, an acyl group optionally having
 substituent(s), an alkoxy carbonyl group optionally
 having substituent(s), a carbamoyl group optionally
 having substituent(s) or a sulfonyl group optionally
 having substituent(s), or


R^2 and R^3 , R^3 and R^4 , or R^4 and R^5

optionally form, together with a carbon atom bonded thereto, a ring optionally having substituent(s),




is an aromatic heterocycle optionally having substituent(s), wherein the aromatic heterocycle is selected from the group consisting of a pyridine ring, pyrimidine ring, a pyridazine ring, a pyrazine ring, a thiophene ring, a furan ring, a pyrrole ring, an imidazole ring, a pyrazole ring, a thiazole ring, an oxazole ring and an isoxazole ring; and the substituent(s) are selected from the group consisting of a halogen atom, an alkyl group optionally having substituent(s), an aryl group optionally having substituent(s), an alkoxyl group optionally having substituent(s), an aryloxy group optionally having substituent(s), an acyloxy group optionally having substituent(s), an alkylthio group optionally having

substituent(s), an arylthio group optionally having
 substituent(s), an acylthio group optionally having
 substituent(s), a protected amino group optionally
 having substituent(s), a nitro group, a cyano group, an
 acyl group optionally having substituent(s), an
 alkoxycarbonyl group optionally having substituent(s), a
 carbamoyl group optionally having substituent(s) and a sulfonyl group optionally
 having substituent(s),
 which comprises reacting a 2-sulfonylpyridine derivative compound represented by the
 formula (I)




wherein

R¹ is an alkyl group optionally having substituent(s) or an aryl group optionally
 having substituent(s), and R², R³, R⁴ and R⁵ are as defined above,
 with an organometallic compound represented by the formula (II')




wherein

M is an atom of an element belonging to Group 1, Group 2, Group 12 or Group 13 of the
 periodic table except a hydrogen atom, and



Claim 2 (Previously Presented): The production method of claim 1, wherein Y¹ is bonded to Y², which X⁴ adjacent to X³ bonded to Y¹ has, to form a double bond, and at least one of Y³ bonded to said X³ and Y⁴ bonded to said X⁴ is an alkyl group optionally having substituent(s).

Claim 3 (Currently Amended): A production method of a pyridine compound ~~derivative~~ having a substituent having a heterocyclic structure at the 2-position, which is represented by the formula (III)



wherein

R², R³, R⁴ and R⁵

are each a hydrogen atom, a halogen atom, an alkyl group optionally having substituent(s), an aryl group optionally having substituent(s), an alkoxyl group optionally having substituent(s), an aryloxy group optionally having substituent(s), an acyloxy group optionally having substituent(s), an alkylthio group optionally having substituent(s), an arylthio group optionally having substituent(s), an acylthio group optionally having

substituent(s), a protected amino group optionally having substituent(s), a nitro group, a cyano group, an acyl group optionally having substituent(s), an alkoxycarbonyl group optionally having substituent(s), a carbamoyl group optionally having substituent(s) or a sulfonyl group optionally having substituent(s), or

R^2 and R^3 , R^3 and R^4 , or R^4 and R^5

optionally form, together with a carbon atom bonded thereto, a ring optionally having substituent(s),

m and n

are each an integer of not less than 1, wherein $m+n=3$ to 8,

C^1 is a carbon atom,

Z is a hydrogen atom, an alkyl group optionally having substituent(s) or an aryl group optionally having substituent(s),

X^1 is a carbon atom, CH, an oxygen atom, a nitrogen atom or a sulfur atom, and


X^2 is a carbon atom, CH, an oxygen atom, a nitrogen atom or a sulfur atom,

wherein at least one of X^1 and X^2 is an oxygen atom, a nitrogen atom or a sulfur atom, when X^1 or X^2 is a carbon atom, CH or a nitrogen atom, Y^1 and Y^2 are each a hydrogen atom, a halogen atom, an alkyl group optionally having substituent(s), an aryl group optionally having substituent(s), an alkoxyl group optionally having substituent(s), an aryloxy group optionally having substituent(s), an acyloxy group optionally having substituent(s), an alkylthio group optionally having substituent(s), an arylthio group optionally having substituent(s), an acylthio group optionally having substituent(s), a protected amino group optionally having substituent(s), a nitro group, a cyano group, an acyl group optionally having substituent(s), an alkoxycarbonyl group optionally having substituent(s), a carbamoyl group optionally having substituent(s) or a sulfonyl group optionally having substituent(s), and

Y^1, Y^2 or Z

is optionally bonded to Y^1 or Y^2 , which X^1 or X^2 adjacent to X^1, X^2 or C^1 bonded to Y^1, Y^2 or Z has, to form a double bond or a ring structure, or when X^1 or X^2 is a carbon atom, Y^1 or Y^2 shows an oxygen atom and is optionally bonded to X^1 or X^2 via a double bond,


which comprises reacting a 2-sulfonylpyridine compound derivative represented by the formula (I)



wherein

R^1 is an alkyl group optionally having substituent(s) or an aryl group optionally having substituent(s), and

with an organometallic compound represented by the formula (II)



wherein

M is an atom of an element belonging to Group 1, Group 2, Group 12 or Group 13 of the periodic table except a hydrogen atom.

Claim 4 (Previously Presented): The production method of claim 3, wherein the organometallic compound (II) has an aromatic heterocycle.

Claim 5 (Previously Presented): The production method of claim 4, wherein the aromatic heterocycle is a pyridine ring, a pyrimidine ring, a pyridazine ring, a pyrazine ring, a thiophene ring, a furan ring, a pyrrole ring, an imidazole ring, a pyrazole ring, a thiazole ring, an oxazole ring or an isoxazole ring.

Claim 6 (Previously Presented): The production method of claim 3, wherein, in the formula (II), M is a lithium atom, a sodium atom, a potassium atom, a magnesium atom, a calcium atom, a zinc atom, a boron atom or an aluminum atom.

Claim 7 (Previously Presented): The production method of claim 3, wherein, in the formula (II), M is a lithium atom or a magnesium atom.

Claim 8 (Previously Presented): The production method of claim 5, wherein the aromatic heterocyclic is a pyridine ring.

Claim 9 (Previously Presented): The production method of claim 6, wherein the aromatic heterocyclic is a pyridine ring.

Claim 10 (Previously Presented): The production method of claim 7, wherein the aromatic heterocyclic is a pyridine ring.